

## STUDIES ON DIFFERENT STOCKING DENSITIES ON PERFORMANCE AND ECONOMICS OF BROILER CHICKEN

**R. D. MAREWAD, M. V. DHUMAL & M. G. NIKAM**

*Department of Poultry Science, College of Veterinary and Animal Science Parbhani,  
Maharashtra Animal and Fishery Science University, Nagpur, Maharashtra, India*

### ABSTRACT

*In order to evaluate the effect of different stocking on performance, litter quality and economics of broiler chicken, an experiment was carried out on by using 400 day old straight run chicks. The experiment was accomplished in completed randomised design including 5 treatments with four replicate and 25 birds per replicate. Treatment group A control with 1.0 sq.ft.area per bird, B with 1.2, C with 1.4 and D with 1.6 sq.ft.area per bird during finisher stage. The results revealed non-significant findings with cumulative body weight gain feed consumption, feed conversion ratio and mortality pattern. Highly significant differences were observed for litter moisture content among various treatment groups. There were non significant differences in weekly pH of litter material. The net profits per Kg live weight, the net profit per square feet area, broiler Performance Efficiency Index (BPEI) and European Productivity Index (EPI) were highest for treatment group A floor space @ 1 sq.ft/bird. Thus the present investigation concluded that increasing stocking density with decreasing floor space resulted into the best broiler performance.*

**KEYWORDS:** *Stocking Density, Floor Space, Broiler Performance, Economics, Litter Moisture and Litter PH*

**Received:** Oct 28, 2016; **Accepted:** Nov 08, 2016; **Published:** Nov 22, 2016; **Paper Id.:** IJASRDEC201648

### INTRODUCTION

The proper stock density is essential for successful broiler production, profitability and welfare point of view. Assessing accurate stocking density, factors such as climate, housing types, ventilation systems, processing weight and welfare regulations must be taken into account. Keeping less birds in the flock is one approach to maintain optimum density.

The optimum density is essential for better performance, however, only providing less space is not a alternative. Improved environmental factors along with space is utmost important. Many factors are responsible for deciding the optimum density like bird size, feeder space, drinker space, house dimension, nutrition, breed, birds welfare, performance, economics. The maximum broiler production per kilogram per square feet area production is achieved by providing optimum density and losses due to overcrowding are prevented. The stocking density is measured is recorded by number of birds per unit area or space per birds. Currently many companies /organizations calculate stocking density by the kilogram. However, the stocking density is as bird weight per unit area. Many recent studies revealed that improved strains of broilers perform better when more space is provided. However, the farmers are not able to afford more than one square area due to unsatisfactory return.

Commercial broiler owners rear the birds in high stocking density for achieving high profitability. However, this may results in higher mortality and lower weight gain. These adverse effects are due to lower air

flow, accumulation of ammonia, carbon monoxide etc. These effects are more pronounced in severe hot humid climatic. Hence, standard stocking density guidelines are needed for recent broiler strains.

The effects of high stocking density on body performance, mortality and some economic indexes of broilers reported to increase ammonia production, foot pad lesions, litter moisture, locomotion, heat stress and preening (8). Stocking density research reported in literature is limited on production parameters, meat yield, and litter quality of broilers grown to body weight of 2.5kg.

From the point, fast growing new strain of broiler and development of new standards, the present experiment was carried on effect of different stocking density on performance and economics of broiler chicken.

## MATERIAL AND METHODS

### Housing and Experimental Design

The experiment was carried out on 400 day old Vencobb-400 straight run commercial broiler chicks for a period of 42 days (6 weeks) from 14<sup>th</sup> January 2015 to 26<sup>th</sup> February 2015 in the Department of Poultry Science, College of Veterinary and Animal Sciences, MAFSU, Parbhani. The experimental design used in the present study for housing the broilers is presented in Table 1 and 2. However the details of percent ingredient composition has been given in Table 3

**Table 1: Experimental Design Used for Housing of Broilers from 0 to 21<sup>st</sup> Days**

| Sr. No | Groups/Treatment | 0 to 7 Days<br>(20% Space) | 8 to 14 Days<br>(40% Space) | 15 to 21 Days<br>(40% Space) |
|--------|------------------|----------------------------|-----------------------------|------------------------------|
| 1      | A                | 0.20 sq. ft.               | 0.30 sq. ft.                | 0.40 sq. ft.                 |
| 2      | B                | 0.24 sq. ft.               | 0.36 sq. ft.                | 0.48 sq. ft.                 |
| 3      | C                | 0.28 sq. ft.               | 0.42 sq. ft.                | 0.56 sq. ft.                 |
| 4      | D                | 0.32 sq. ft.               | 0.48 sq. ft.                | 0.64 sq. ft.                 |

**Table 2: Experimental Design Used for Housing of Broilers from 22<sup>nd</sup> to 42<sup>nd</sup> Days**

| Treatment Group              | Treatment Group Details                                 | No. of Birds/Pen/<br>Replication | No. of<br>Replication | Total<br>Number<br>of Birds |
|------------------------------|---|----------------------------------|-----------------------|-----------------------------|
| A                            | Control group; 1 sq.ft. Floor space per bird            | 25                               | 4                     | 100                         |
| B                            | Increase Floor space by 20% to reach 1.2 sq.ft.per bird | 25                               | 4                     | 100                         |
| C                            | Increase Floor space by 40% to reach 1.4 sq.ft.per bird | 25                               | 4                     | 100                         |
| D                            | Increase Floor space by 60% to reach 1.6 sq.ft.per bird | 25                               | 4                     | 100                         |
| <b>Total Number of Birds</b> |   |                                  |                       | <b>400</b>                  |

The birds were randomly distributed with maintaining 50:50 sex ratios to form the experimental groups. The floor space was provided for all the treatment groups excluding feeding and drinking space. The same number of feeders and drinkers for all the experimental treatment groups were provided.

**Table 3: Percent Ingredient and Nutrients Composition of Feed Starter, Grower and Finisher Ration of Broiler Chicken**

| Feed Ingredients | Starter<br>(0-10d) | Grower<br>(11-21d) | Finisher<br>(22-42d) |
|------------------|--------------------|--------------------|----------------------|
| Maize            | 53.2               | 54.0               | 58.5                 |
| Vegetable oil    | 3.0                | 4.2                | 4.9                  |

| Table 3: Contd.,                        |          |          |          |
|---|----------|----------|----------|
| Soya-bean meal                          | 41.0     | 39.0     | 33.8     |
| Dicalcium phosphate                     | 1.5      | 1.5      | 1.5      |
| Limestone powder                        | 1.0      | 1.0      | 1.0      |
| Salt                                    | 0.3      | 0.3      | 0.3      |
| Total                                   | 100      | 100      | 100      |
| <b>Supplements/Additives (g/100kgs)</b> |          |          |          |
| *Mineral mixture                        | 300      | 300      | 300      |
| **Vitamin mixture                       | 150      | 150      | 150      |
| Methionine                              | 180      | 190      | 160      |
| Lysine                                  | 170      | 130      | 100      |
| Choline chloride                        | 60       | 60       | 60       |
| Nutritional content (calculated)        |          |          |          |
| Crude protein (%)                       | 23.08    | 22.24    | 20.22    |
| Metabolizable energy (Kcal/kg)          | 3011.16  | 3100     | 3202.22  |
| E:P ratio                               | 130.46:1 | 139.38:1 | 158.37:1 |

\*Mineral Mixture: - Copper, ferrous, zinc, iron.

\*\*Vitamin Mixture: -Vit.A, D, E, K & Vit B Complex (Riboflavin, thiamine, Choline, pantothenic acid, niacin, pyridoxine, biotin, cynocobalamine).

The feed ingredients used in the present experiment were purchased from local market and rations were prepared as per (6) at Feed Mixing Plant, College of Veterinary and Animal Sciences, MAFSU, Parbhani. (Table 3)

### Feeding and Watering Schedules

Weighed quantity of feed was offered to each replicate of treatment groups. The left over feed was collected and weighed separately on the first day morning of second week and onward of each week to arrive at the actual weekly feed consumption, which was inclusive of feed losses, if any. The birds were offered adlib fresh and clean drinking water throughout the experiment.

### Data Collection

Data were collected on weekly weight changes was determined by weighing the birds on weekly basis and weight gain was calculated by subtracting the weight of the previous week from that of the current week. The feed intake was determined by subtracting the left-over feed from the feed offered, while feed conversion ratio was calculated as average feed intake divided by average /weight gain. The mortality was recorded daily and the weights of all the dead birds were taken in order to minimize an error in feed conversion ratio. It was expressed as percentage mortality at the end of the experiment for corresponding treatment group.

### Moisture Percentage of Litter Material

The litter samples were collected from four locations within each pen. (Four peripheral, equidistant from each pen corner) and thoroughly mixed to obtain material representative of the entire pen. At least 200 gram of litter was placed in a plastic container and a subsample was taken for further analysis at the laboratory. The litter moisture samples were collected, mixing and obtaining a 100 gram litter subsample, placed in 15x30 cm tray and oven-dried for 48 hours at 60C. The percentage of moisture was calculated by using the weight loss between initial and dried litter (37) by using following formula.

|             |                         |       |
|-------------|-------------------------|-------|
| Moisture% = | (initial wt – dried wt) | X 100 |
|             | initial wt              |       |

Where Wt = weight

### Estimation of pH of Litter Material

The upper 10 cm of the litter was collected at each sample position for determination of pH. The pH of each sample was measured after litter samples of nearly 5 gram were suspended for 30 min in 25 ml of distilled water and stirred for 5 min using a pH meter (Mettler Toledo, GmbH, Switzerland, 25).

### Economics of Broiler Production

The economics of broiler production was calculated by considering the cost of chick, total feed, litter, vaccine, medicine. However, the labour charges are not considered because the experiment was carried out by the student. The gross profit of broiler production per bird was calculated by subtracting the cost of production per bird from market price fetched on live weight basis.

### Broiler Performance Efficiency Index

The index took into account total salable weight, feed efficiency as well as livability.

|        |                                 |      |
|--------|---------------------------------|------|
| BPEI = | Total Saleble Live Weight (kg)  | X100 |
|        | No. of Chicks x Feed Efficiency |      |

### European Production Index

The index took into account average live weight, safety as well as feed consumption and total growing days.

|       |   |      |
|-------|---|------|
| EPI = | Average Live Weight (kg) x Safety %                   | X100 |
|       | Feed Consumption Per Kg of Gain x Term of Growing day |      |

### Statistical Analysis

All the generated data were subjected to statistical analysis (ANOVA) by using Complete Randomized Design (34). The treatment means were compared by critical differences (CD) and Analysis of Variance.

## RESULTS AND DISCUSSIONS

### Cumulative Weight Gain

The results of weight gain are presented in Table 4 & 5. The results showed that there were no significant differences in weight gain between the treatments in all ages. Non-significant results in regard to body weight gain of broiler in stocking density of 1, 1.2, 1.4 and 1.6 square feet per bird indicated increased creation of metabolic energy. An young broiler chickens can use for rapid growth, resulted increase in feed consumption and metabolism led to increased creation of heat (38). The weight gain observed in the present study with different floor space revealed the statistically non-significant differences amongst the treatment groups which may indicative of the fact that the increasing floor space per bird may not provide the required comfort zone for gaining the body weight with increasing floor space. However, the critical analysis of the numerical values of the cumulative weekly weight gain revealed the universally accepted fact and phenomenon of increasing stocking density with decreasing floor space (from 1.6 to 1.0 sq.ft. floor space per bird) resulted into the best performance in terms of body weight gains in the broilers. The results are similar in body weight gain of birds reared at different density (5) and found no differences in final body weight at 39 days of age with increasing stocking density. Similarly (26) reported that increase in population density in broiler chicken was not significantly influenced on the final mass of broilers. Similar findings were also reported by (19), (36) and (18). However, findings in the present

study are in contrast to the reports of many scientists (14,31,10,15, 31, 21,9 &22).

**Table 4: Weekly Cumulative Weight Gain (g), Feed Consumption, FCR and Mortality at Different Age Groups with Different Stocking Density**

| Age (Weeks)                 | Groups/Treatments      |               |               |                |
|-----------------------------|------------------------|---------------|---------------|----------------|
|                             | Cumulative Weight Gain |               |               |                |
|                             | A (1.0 Sq.ft.)         | B (1.2Sq.ft.) | C (1.4Sq.ft.) | D (1.6 Sq.ft.) |
| I                           | 113.88                 | 117.72        | 114.55        | 122.77         |
| II                          | 355.45                 | 362.72        | 360.18        | 372.00         |
| III                         | 752.84                 | 746.62        | 749.06        | 762.25         |
| IV                          | 1256.73                | 1231.32       | 1234.96       | 1259.76        |
| V                           | 1849.99                | 1772.07       | 1797.21       | 1832.50        |
| VI                          | 2369.77                | 2315.77       | 2333.98       | 2331.92        |
| Cumulative Feed Consumption |                        |               |               |                |
| I                           | 136.63                 | 136.76        | 136.91        | 140.80         |
| II                          | 440.20                 | 440.26        | 440.71        | 445.10         |
| III                         | 985.21                 | 983.29        | 985.65        | 995.70         |
| IV                          | 1769.32                | 1786.96       | 1776.06       | 1800.03        |
| V                           | 2735.81                | 2762.63       | 2749.89       | 2792.76        |
| VI                          | 3756.44                | 3763.17       | 3766.92       | 3805.63        |
| Cumulative FCR              |                        |               |               |                |
| I                           | 1.20                   | 1.16          | 1.20          | 1.17           |
| II                          | 1.24                   | 1.21          | 1.22          | 1.20           |
| III                         | 1.31                   | 1.32          | 1.32          | 1.31           |
| IV                          | 1.41                   | 1.45          | 1.44          | 1.43           |
| V                           | 1.48                   | 1.56          | 1.53          | 1.52           |
| VI                          | 1.59                   | 1.62          | 1.61          | 1.63           |
| Mortality (%)               |                        |               |               |                |
| Overall Mortality           | 0                      | 0             | 1             | 1              |

**Table 5: Analysis of Variance of Cumulative Weight Gain (g) of Broilers at Different Age Groups with Different Stocking Density**

| Source | DF | Cumulative Weight Gain |            |         | Cumulative Feed Consumption |             |           | Cumulative FCR |       |         |
|--------|----|------------------------|------------|---------|-----------------------------|-------------|-----------|----------------|-------|---------|
|        |    | SS                     | M SS       | F ratio | SS                          | MSS         | F Ratio   | SS             | MSS   | F Ratio |
| Groups | 3  | 10638.846              | 3546.282   | 1.757   | 9172.946                    | 3057.64     | 1.423     | 0.006          | 0.002 | 0.721   |
| Weeks  | 5  | 59070600.              | 11814100.0 | 5853.4  | 159100000.00                | 31820000.00 | 14811.439 | 2.382          | 0.476 | 180.525 |
| Error  | 87 | 175592.415             | 2018.304   |         | 186905.62                   | 2148.341    |           | 0.230          | 0.003 |         |
| Total  | 95 | 59256830.261           |            |         | 159286078.57                |             |           | 2.618          |       |         |

### Cumulative Feed Consumption

The results of cumulative feed consumption are presented in Table 4&5. The results showed that there were no significant differences in feed consumption between the treatments in all ages. Statistically non-significant differences between the treatment groups indicating the fact that stocking density and floor space per bird (1 to 1.6 sq. ft. /bird) did not play any significant role in feed consumption. However, the critical observation of cumulative feed consumption in the present study revealed the fact that increasing floor space or decreasing stocking density increased the cumulative feed consumption. This may lead to the conclusion that increased floor space per bird engaged the bird only on concentrating feeding activity. This might have resulted in numerically increased feed consumption. These results in the present study

were similar to the findings reported by many scientists (28, 24, 36, and 18). However, the present findings were in contrast to the finding (19), (21), (20), (35), they revealed that the feed consumption was influenced ( $P < 0.01$ ) by stocking density.

### **Cumulative Feed Conversion Ratio (FCR)**

The results of cumulative feed consumption are presented in Table 4 & 5. The statistically non-significant differences observed for cumulative feed conversion ratio amongst the different treatment groups provided with different stocking densities/ floor space. The critical observation of the weekly and cumulative feed conversion ratio did not reveal even the numerical differences which may lead to the strong conclusion that the broiler birds (Vencobb-400) must have a very good genetic potentiality of exhibiting most efficient feed conversion ratio ability even though they are subjected to different floor space and stocking densities when other managerial factors were at par. Better feed efficiency in higher stocking densities was established by (14), Similarly, (33) established better feed conversion ratio in all stages of broiler nutrition in higher stocking density. Absence of the effect of stocking density on feed conversion ratio was also concluded by (10). The numerous authors confirm non-significant effect of stocking density on feed conversion ratio. (29, 35, 30, 36 & 18). Whereas, (19), (3), (24) revealed significant influence of stocking density on feed conversion ratio.

### **Mortality**

The results of mortality are presented in Table 4 & 5. Non-significant influence of stocking density revealed that the space provided at 1.0 square feet per bird is optimum. The findings in accordance with present study were reported by (14), (29), (35), (24), & (30). In regard to mortality, majority of authors revealed non-significant differences of different stocking density, although it should not be overlooked that percentage of died broiler birds increases with the increase of stocking density (33), especially in stocking density below 1 square feet. In contrast to present study findings (20) showed significant increase of mortality with the increase of stocking density.

### **Moisture Percentage of Litter Material**

The results of moisture percentage of litter material are presented in Table 7 & 8. Moisture percentage of litter material was significantly ( $P < 0.05$ ) influenced by stocking density in the present study. Significantly higher moisture percentage of litter observed for treatment group A provided with 1 square feet area compared to other treatment groups at the 6<sup>th</sup> week of age. However, the moisture of litter was within ideal limit (20-25 %, 17). If litter moisture exceeds more than 25%, it will promote the proliferation of pathogenic bacteria and moulds and wet litter is also the primary cause of the high level of ammonia. Controlling litter moisture is most important step in avoiding ammonia problems (4). To limit ammonia production, litter moisture below 30% and temperature at the level of birds comfort is essential. In the present study the litter moisture was within optimum condition. The litter that is too dry and dusty can also leads to problems of respiratory diseases. Caked litter correspond to high litter moisture, litter becomes anaerobic. The moisture content of litter exceeding 35% is detrimental and correlated with higher fecal corticosteroid hormone which can increase the mortality. (7). The findings in the present study are in accordance with (10), (33), (29) & (30) revealed that litter moisture content of the groups were significantly affected by stocking density. However, (18) reported that, at 45 days of age, litter moisture did not differ statistically between two densities, in spite of the numerical differences in treatments.

**Table 6: Weekly Moisture Percentage and pH of Litter Material with Different Stocking Density**

| Age (Weeks)              | Treatment Groups                          |               |               |                |
|--------------------------|---|---------------|---------------|----------------|
|                          | A) Moisture Percentage of Litter Material |               |               |                |
|                          | A (1.0 Sq.ft.)                            | B (1.2Sq.ft.) | C (1.4Sq.ft.) | D (1.6 Sq.ft.) |
| I                        | 5.85                                      | 5.59          | 5.77          | 5.57           |
| II                       | 26.25                                     | 32.45         | 28.49         | 26.57          |
| III                      | 32.47                                     | 29.45         | 26.69         | 27.77          |
| IV                       | 26.19                                     | 27.05         | 26.46         | 25.14          |
| V                        | 30.84                                     | 27.18         | 25.94         | 21.67          |
| VI                       | 32.08                                     | 21.49         | 23.55         | 20.90          |
| B) pH of Litter Material |   |               |               |                |
| I                        | 6.23                                      | 6.26          | 6.25          | 6.27           |
| II                       | 5.96                                      | 5.71          | 5.86          | 5.91           |
| III                      | 5.62                                      | 6.41          | 6.29          | 5.88           |
| IV                       | 6.58                                      | 7.07          | 6.96          | 6.67           |
| V                        | 6.96                                      | 6.59          | 6.98          | 6.69           |
| VI                       | 7.07                                      | 6.78          | 6.67          | 6.53           |

**Table 7: Analysis of Variance of Moisture Percentage and pH of Litter Material with Different Stocking Density**

| Source       | DF        | Moisture Percentage of Litter Material |         |          | pH of Litter Material |       |         |
|--------------|-----------|--|---------|----------|-----------------------|-------|---------|
|              |           | SS                                     | M SS    | F ratio  | SS                    | MSS   | F ratio |
| Groups       | 3         | 384.042                                | 128.014 | 7.471**  | 0.728                 | 0.243 | 1.758   |
| Weeks        | 5         | 984.175                                | 196.835 | 11.487** | 17.025                | 3.405 | 24.677  |
| Error        | 87        | 1490.776                               | 17.135  |          | 12.005                | 0.138 |         |
| <b>Total</b> | <b>95</b> | <b>59256830.261</b>                    |         |          | <b>29.758</b>         |       |         |

**pH of Litter Material**

The results of pH of litter material are presented in Table 7 & 8. A non-significant effect of stocking density on pH of litter material in the present study indicated ideal pH. To control ammonia production the litter pH should be below 7.0. In the present study pH of litter was within the ideal and normal limits. (29), (13) and (30) had revealed the significant to highly significant effect of increasing stocking density with the increased pH of litter material. It was in contrast to the present study where non-significant differences between treatment groups of stocking density were observed.

**Economics of Broiler Production**

The economics of broiler production from different treatment groups is presented in Table 9. The cost of day old chicks, feed, medication, vaccination, litter and other overheads were considered while calculating the cost of production. However, the costs of labour were not considered in calculating the cost of production of the broilers as this experiment being a postgraduate research work.

**Table 8: Economics of Broiler Production at Different Stocking Density**

| Sr. No. | Particulars                | Treatment Groups |               |               |               |
|---------|----------------------------|------------------|---------------|---------------|---------------|
|         |                            | A (1.0Sq.ft.)    | B (1.2Sq.ft.) | C (1.4Sq.ft.) | D (1.6Sq.ft.) |
| 1       | Cost of day old chick (Rs) | 20               | 20            | 20            | 20            |
| 2       | Feed consumption (g)       |                  |               |               |               |
| i)      | Prestarter                 | 501.44           | 503.17        | 500.92        | 505.63        |
| ii)     | Starter                    | 1002.50          | 1005.20       | 1006.50       | 1010.35       |

| Table 8: Contd., |   |                |                |                |                |
|------------------|---|----------------|----------------|----------------|----------------|
| ii)              | Finisher  | 2252.50        | 2254.80        | 2259.50        | 2289.65        |
|                  | <b>Total</b>                                      | <b>3756.44</b> | <b>3763.17</b> | <b>3766.92</b> | <b>3805.63</b> |
| 3                | Rate of feed (Rs/kg)                              |                |                |                |                |
| i)               | Prestarter  | 26.32          | 26.32          | 26.32          | 26.32          |
| ii)              | Starter   | 26.44          | 26.44          | 26.44          | 26.44          |
| iii)             | Finisher  | 25.59          | 25.59          | 25.59          | 25.59          |
| 4                | Cost of feed consumed (per bird Rs.)              |                |                |                |                |
| i)               | Prestarter  | 13.19          | 13.24          | 13.18          | 13.30          |
| ii)              | Starter   | 26.50          | 26.57          | 26.61          | 26.71          |
| iii)             | Finisher  | 57.64          | 57.70          | 57.82          | 58.59          |
|                  | <b>Total Cost of Feed Consumed per Bird (Rs.)</b> | <b>97.33</b>   | <b>97.51</b>   | <b>97.61</b>   | <b>98.60</b>   |
| 5                | Miscellaneous cost* (Rs)                          | 5              | 5              | 5              | 5              |
| 6                | Total cost of production (1+4+5)                  | 122.33         | 122.51         | 122.61         | 123.60         |
| 7                | Average live weight (g)                           | 2405.87        | 2351.76        | 2370.03        | 2367.92        |
| 8                | Return obtained @ Rs. 58 per kg live weight       | 139.54         | 136.40         | 137.46         | 137.34         |
| 9                | Net Profit/Bird (Rs.)                             | 17.21          | 13.89          | 14.85          | 13.74          |
| 10               | Net profit/ kg (Rs)                               | 7.15           | 5.90           | 6.26           | 5.80           |
| 11               | Net profit per sq.ft area                         | 17.21          | 11.58          | 10.61          | 8.59           |
| 12               | Broiler performance efficiency index(BPEI)        | 175.61         | 169.31         | 169.17         | 170.37         |
| 13               | European Productivity Index (EPI)                 | 418.12         | 403.13         | 402.78         | 405.63         |

Significant findings in the present study for net profit per bird, net profit per kg, net profit per square feet area, broiler performance efficiency index (B.P.E.I.) and European productivity index (E.P.I.) for treatment group A compared to rest of the treatment groups concluded that higher live body weight in broiler maintained under less floor space increased the profitability. The research findings in the present study were in accordance with (23), they revealed that reduction in final stocking density caused drop in net income from broiler production. The profitability enhanced with increased stocking density (30). It may be due to reduction of fixed cost and more kilogram production of broiler per area.

## CONCLUSIONS

Based on present study, it is concluded that a limitation of floor space for broiler farming should go for high density (1 square feet area per bird) with precise system for ventilation that could effectively used for soothing the bird's micro climate. When the poultry farmer's focus is on body size and body weight rather than number and keeping birds more than 45 days they should opt low bird density (1.2 square feet area per bird). The economics of broiler observed in present study concluded that direct and proportionate correlation of increasing floor space with decreasing profit in all aspects of profit calculation which may strongly lead to the conclusion that 1 sq. ft./ bird floor space is the most ideal and realistic in all respects of broiler production.

## ACKNOWLEDGEMENTS

The authors are highly grateful to S B Majee, Associate Dean, College of Veterinary and Animal Sciences, MAFSU, Parbhani for providing necessary facilities and help at various stages of the experiment.



## REFERENCES

1. Abudabos Alaeldein M., Emad M. Samarag, O. S. Hussein, Muith Q. Al-Ghadi, Raed M. Al-Atiyat 2013. Impacts of stocking density on the performance and welfare of broiler chickens. *Italian Journal of Animal Sciences*; 12e11
2. Ali M. I., S. A. Azmal, A. Ali and M. O. Faruque 2012. Effect of density and flock size on growth performance of native chicken. *J.Bangladesh agric. univ.* 10 (1): 55-59.
3. Asaniyan E. K 2014. Growth performance, feed intake and slaughter characteristics of broiler chicken under three different stocking densities. *J. Anim Prod Adv*, 4 (2): 348-354
4. Atilgan A., Coskan A., Oz. H., Isler E 2010. Etlik pilic kumesinde kis doneminde amonyak gaz duzeyinin vakum sistemi ile azaltılması. *Kafkas Univ Vet Fak Derg*, 16, 257-262.
5. Buijs S, L. Keeling, S. Rettenbacher, E. Van Poucke, and F. A. M. Tuytens 2009. Stocking density effects on broiler welfare: Identifying sensitive ranges for different indicators. *Poultry science* 88:1536-1543.
6. Bureau of Indian standards 2007. Indian standards. *Poultry feed specification (fifth revision)* page 3-5.
7. Butcher G. D., Miles R. D. (2009). Causes and prevention of wet litter in broiler houses, University of Florida IFAS Extension VM 99.
8. Cravener, T. L., W. B. Roush, and M. M. Mashaly.1992. Broiler production under varying population densities. *Poult. Sci.* 71:427-433.
9. Das, H and Lacin, E 2014. The effect of different photoperiods and stocking densities on fattening performance, carcass and some stress parameters in broilers. *Israel Journal of Veterinary Medicine*. 69 4.
10. Dozier W. A, III, J. P. Thaxton, S. L. Branton, G. W. Morgan, D. M. Miles, W. B. Roush, B. D. Lott and Y. Vizzier Thaxton 2005. Stocking density effects on growth performance and processing yields of heavy broilers. *Poultry Science* 84:1332-1338.
11. Estevez, I. 2007. Density allowances for broilers: where to set the limits? *Poult. Sci.* 86.1265-1272.
12. Fairchild Brian D.2005. Broiler tip. The University of Georgia. Cooperative Extension Service. college of Agricultural and environmental Sciences/ Athens, Georgia 30602-4356.
13. Farhadi, Danial and Hosseini, Seyed Majid. 2014. Comparison of broiler performance in two conventional and environmentally controlled conditions modern broiler houses in tropics. *Global Journal of Animal Scientific Research*. 2 (3):190-196.
14. Feddes, J. J. R., Emmanuel, E. J. and Zuidhof. M. J. 2002. Broiler performance, body weight variance, feed and water intake, and carcass quality at different stocking densities. *Poul. Sci.* 81: 774-779.
15. Ghorpade Prerna, Barmase B. S., Kuralkar S. V., Ambulkar D. R.2008. Effect of different housing system and stocking density on growth performance of broilers. *Indian J. of Poultry sci.* 43:116-119.
16. Ghosh Sarbaswarup, Debasis Majumder and Rupak Goswami 2012. Broiler performance at different stocking density. *Indian J. Anim. Res.* 46 (4):381-384
17. Glebocka K. 2008. Gut health is a critical factor for litter quality. *World Poultry*. 24:12-13.
18. Gopinger Edenilse, Vaidir Silveira de Avila, Dani Perondi, Aiane Aparecida da Silva Catalan, Everton Luis Krabbe and Victor Fernando Buttow Roll 2015. Performance, carcass characteristics and litter moisture in broilers housed at two densities. *Acta Scientiarum. Animal sciences*. Maringa. 37. 1, 35-39.

19. Guardia S., B. Konsak, S. Combes, F. Levenez, L. Cauquil, J. F. Guillot, C. Moreau- Vauzelle, M. Lessire, H. Juin, and I. Gabriel 2011. Effects of stocking density on the growth performance and digestive microbiota of broiler Chickens. *Poultry science* 90; 1878-1889.
20. Hall A. L. 2001. The effect of stocking density on the welfare and behaviors of broiler chickens reared commercially. *Anim. Welfare*, 10, 23-40.
21. Hassenein H. H. M.2011. Growth performance and carcass yield of broilers as affected by stocking density and enzymatic growth promoters. *Asian journal of poultry sciences* ISSN 1819-3609.
22. Jayalakshmi T., R. Kumararaj, T. Sivakumar, Thanga Thamil Vanan and D. Thiagarajan 2009. Influence of stocking densities on litter moisture, microbial load, air ammonia concentration and broiler performance. *Tamilnadu j. veterinary & animal sciences*. 5 (3) 80-86.
23. Katarzyna Utnik-Banas, Janusz Zmija, Elzbieta Sowula-skrzynska 2014. Economic aspects of reducing stocking density in broiler chicken production using the example of farms in southern Poland. *Ann. anim. Sci.*, 14:3:663-671. DOI: 10.2478/aoas-2014-0029.
24. Mehmood S, A W Sahota, M Akram, K. Javed J. Hussain, M. S. Shaheen, Y. Abbas, A. S. Jatoi and A. Iqbal 2014. Growth performance and economic appraisal of phase feeding at different stocking densities in sexed broilers. *The J. of Animal & Plant Sciences*, 24(3):714-721.
25. Miles DM, Owens PR, and Rowe DE 2006. Spatial variability of litter gaseous flux within a commercial broiler house: ammonia, nitrous oxide, carbon dioxide, and methane. *Poultry Sci.* 85:167-172.
26. Mitrovic S., V. Dermanovic, Mihailo Radivojevic, Zoran Rajic, Dragic Zivkovic, Dorde Ostojic and Nikola Filipovic 2010. The influence of population density and duration of breeding on broiler chickens productivity and profitability. *African J. of Biotechnology*. 9(28), 4486-4490.
27. Mortari, A. C., Rosa, A. P., Zanella, I., Neto, C. B., Visentin P. R. and Brites, L. B. P. : 2002. Performance of broilers reared in different population density, in winter, in South Brazil. *Ciencias Rural*. 32: 493-497.
28. Nogueira WCL; Velasquez PAT; Furian RL; M Macari 2013. Effect of dietary energy and stocking density on the performance and sensible heat loss of broilers reared under tropical winter conditions. *Department of Animal Morphology and physiology. Faculdade de Ciencias Agrarias e Veterias – UNESP. Jaboticabal campus, Sao Paulo, Brazil*.
29. Petek, M., Recep Cibik, Huseyin Yildiz, Fisun AK Sonat, Serife Sule Gezen, Abdulkadir Orman, Cenk Aydin 2010. The influence of different lighting programs, stocking densities and litter amounts on the welfare and productivity traits of a commercial broiler line. *Veterinarija ir Zootechnica (vet med zoot)*.T.51 (73).
30. Petek, M., Hakan Ustuner, Derya Yesilbag 2014. Effects of stocking density and litter type on litter quality and growth performance broiler chicken. *Kafkas Univ Vet Fak Derg* 20(5):743-748.
31. Ravindran, V., Donald V., Thomas, David G., Thomas and Patrick C. H. Morel 2005. Performance and welfare of broilers as affected by stocking density and zinc bacitracin supplementation. *Institute of food, nutrition and human health, Massey university Palmerston north, New Zealand*.
32. Silas, Abel F. A. Adeyemi Olajide Ayorinde, Erubberine Daisy, Sogunle Olajide Mark, Oluwole Oluwatoyin Bolanale, Elemo Gloria Nwakaegho 2014. Effect of stocking density and quantitative feed restriction on growth performance, digestibility, hematological characteristics and cost of starting broiler chicks. *Journal of Animal Health and production* 2 (4):60-64.

33. Skrbic. Z., Pavlovski, M. Lukic 2009. Stocking density -factor of production performance, quality and broiler welfare. *Biotechnology in Animal Husbandry* 25 (5-6). p 359-372.
34. Snedecor. W. and Cochran, W. G. 2002. *Statistical Methods*, 6<sup>th</sup> Ed., Oxford and IBH publishing Co., Calcutta, Bombay and New Delhi, pp 168-181.
35. Uzum, M. H., H. D. Oral Toplu 2013. Effects of stocking density and feed restriction on performance, carcass, meat quality characteristics and some stress parameters in broilers under heat stress. *Revue Med. Vet.* 164, (12), 546-554.
36. Wang, Bo. Zhizhi Min, Jiyanmin Yuan, Bingkun Zhang and Yuming Guo 2014. Effects of dietary tryptophan and stocking density on the performance, meat quality, and metabolic status of broilers. *Journal of Animal Science and Biotechnology* 5:44.
37. Willis WL. Murray C., Talbott C: (1997). Evaluation of leaves as a litter material. *Poultry Sci*, 76.1138-1140
38. Yadgari. L. R. Kinereich, S. Druyan, & A. Cahaner, 2006. The effects of stocking density under hot condition on growth, meat yield and meat quality of featherless and feathered broilers. *Proc. 12<sup>th</sup> Eur. Conf. World's Poultry Science Journal*, 62 (supplement). 603-604.

